

THE LARGE-DISDROMETER EXPERIMENT (LDE) & UCLM'S X-BAND RADAR (T-REX) STATUS

Francisco J. Tapiador

+

Andrés Navarro, Raúl Moreno, Alfonso Jiménez, Cecilia Marcos

[thanks to Walt Petersen, Mathew Schwaller, David Wolff,
Ziad S. Haddad, Chris Kummerow, Ali Tokay, and Ramesh Kakar]


UCLM's Disdrometers


2 Thies


16 Parsivel-1

8 Parsivel-2


Several experiments








Thies disdrometers on top of a
100m meteo tower and at 5m



CIBA

Some experiments:

- Comparing 100m-5m measurements for the raining events
- Just a few episodes
- Database still to be exploited




2009 PMM Science Team Meeting, Salt Lake City, UT

The UCLM D-Array


Made of 9 times this dual setup (18 Parsivels)








First setup, 2009




F. J. Tapiador; R. Checa; M. de Castro. 2010.
An experiment to measure the spatial variability of rain
drop size distribution using sixteen laser disdrometers.
GEOPHYSICAL RESEARCH LETTERS. 37


Reflectivity (Z)



F. J. Tapiador; R. Checa; M. de Castro. 2010.
An experiment to measure the spatial variability of rain
drop size distribution using sixteen laser disdrometers.
GEOPHYSICAL RESEARCH LETTERS. 37







outlier



Reflectivity (Z)

Earth and Space Sciences (ess) Group, UCLM, Toledo, Spain





Parsivel¹ + Parsivel²

2015

Parsivel² improves over Parsivel¹

[more sensitive to small drops]


The 'Large Disdrometer Experiment'

How important is the small catching area of disdrometers? Large biases?


How large should it be?



LDE
2011




Experimental layout







The catching area is larger:
increased probability of having
rare events (large drops)

Assumption: Estimates are going
to be closer to the truth than
those from just one disdrometer

The synthetic LD acts here as a
reference: our “true” estimate of
the DSD






One disdrometer




Large-Disdrometer
(LD) equivalent area



What if we have one?


That would give us a different
answer from the LD



What if we have another one?

That would also give us a different answer


Different from the first, and also from the LD estimate



What if we have another one?

That would also give us a different answer

Different from the first, and from the LD estimate




For $r=1$, we have 16 arrangements

Cause we have $\binom{n}{r}$ ("n choose r")

arrangements of r disdrometers
taken from a set of n disdrometers


(also noted as C_r^n)



What if we have two?

That would give us a different answer

Different from just one, and different from the LD-reference




[illegible]

Same thing for other arrangements

What if we have two?

That would give us a different answer

We have $16!/(14! \cdot 2!) = 120$ combinations




What if we have six?

That would indeed give us a different answer


We have now 8008 combinations

But this answer is likely to be closest to the 'truth' than having just two




In total, we calculated 16k different estimates

We **didn't** randomly selected a setup, we calculated all the possible combinations.




Of course we did not just aggregate the results from individual disdrometers

We didn't do this:




Of course we did not just aggregate the results from individual disdrometers


We didn't do this:




We rather input the processing software (Tokay's) with the data of the **union** of data from several disdrometers then calculate the mean values over the combinations, over time



Catching area



Catching area




A single disdrometer almost certainly:

- Underestimates Z
- Overestimates R
- Overestimates D_{mass}
- Underestimates D_{max}



This is something we should be aware of when considering the values given by a single instrument

The combined area of 7 disdrometers seems enough for this case










There is of course other
sources of error/uncertainty

Thus for instance in calculating
radiometric quantities








[spherical drops; drop size is exaggerated in these plots]



[spherical drops; drop size is exaggerated in these plots]

Ensemble of 50
arrangements
a 1 m^3 space







**T-matrix multiple scattering
calculations for a variety of
arrangements, frequencies and
temperatures**

- **Complex refractive index
depends on temperature,
(salinity) and frequency**
- **Length scale factor depends on
frequency**


X-band; 5 °C

**Spread of radiometric quantities
within the ensemble (same DSD)**




X-band; 5 °C

CASE A



CASE B



UCLM's T-Rex

A dual polarization X-band radar



Transmitter (Magnetron type)

featuring:

- Frequency range: **9.375 GHz (X-Band)**
- Nominal RF Peak Power > 55 kW
- Pulse modes: 3
- Pulse Width: 0.33 μ s - 3.3 μ s

SELEX METEOR 60DX TYPE



RF Receiver Unit

featuring:

- Frequency range 9.3 - 9.5 GHz (X-Band)
- Noise figure < 2.5 dB
- RX total dynamic range > 90 dB
- Dual polarization, H + V simultaneous



[Blame Walt]




T-Rex at Wallops

New (received December 2015)

Now at Wallops

We hope we can start
measuring by next week

ICE-POP 2018 [slide from G. Lee]



Thanks